

FORM PTO-1449 (modified)
To: U.S. Department of Commerce
(PW FORM PAT-1449)
Patent and Trademark Office

Atty. Dkt. No.: 068800-0324485

**INFORMATION DISCLOSURE STATEMENT
BY APPLICANT**

Applicants: PEPYS et al.

Application No. : 10/559,814

Filing Date:

Date: June 21, 2007

Page

1

of

5

Examiner: Ruixiang Li Group Art Unit: 1646

U.S. PATENT DOCUMENTS

| Examiner's Initials* | | Document Number | Date MM/YYYY | Name (Family Name of First Inventor) | Class | Sub Class | Filing Date (if appropriate) |
|-------------------------|----|--------------------|-----------------|---|-------|--------------|---------------------------------|
| | AR | 6,365,570 | 04/2002 | Van Kessel et al. | 514 | 8 | |
| | BR | 6,126,918 | 10/2000 | Pepys et al. | 424 | 9.1 | |
| | CR | 7,045,499 B2 | 05/2006 | Pepys | 514 | 2 | |
| | DR | 4,783,480 | 11/1988 | Wakatsuka et al. | 514 | 423 | |
| | ER | 4,895,872 | 01/1990 | Nitecki et al. | 514 | 546 | |
| | FR | 6,103,910 | 08/2000 | Hertel et al. | 548 | 523 | |

FOREIGN PATENT DOCUMENTS

| | | Document Number | Date MM/YYYY | Country | Inventor Name | English Abstract | | Translation Readily Available | |
|--|----|--------------------|-----------------|---------|-----------------|---------------------|----|-------------------------------------|----|
| | | | | | | Enclosed | No | Enclosed | No |
| | GR | EP 0915088 A1 | 05/1999 | EPO | Hertel et al. | Yes | | Yes | |
| | HR | WO 95/05394 | 02/1995 | WIPO | Pepys et al. | Yes | | Yes | |
| | IR | WO 97/46098 | 12/1997 | WIPO | Theodore et al. | Yes | | Yes | |
| | JR | WO 98/50420 | 11/1998 | WIPO | Adang et al. | Yes | | Yes | |
| | KR | WO 03/013508 | 02/2003 | WIPO | Pepys | Yes | | Yes | |

OTHER (Including in this order Author, Title, Periodical Name, Pertinent Pages, Date, etc.)

| | | |
|--|----|--|
| | LR | |
| | MR | |
| | NR | |
| | OR | |
| | PR | |
| | QR | |
| | RR | |
| | SR | |
| | TR | |

Examiner /Ruixiang Li/

Date Considered: 02/27/2009

*EXAMINER: Initial if citation considered, whether or not citation is in conformance with MPEP § 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

ALL REFERENCES CONSIDERED EXCEPT WHERE LINED THROUGH. /R.L./

Atty. Dkt. No.: 068800-0324485

**INFORMATION DISCLOSURE STATEMENT
 BY APPLICANT**

Applicant: PEPYS

Application No. : 10/559,814

Filing Date:

Date: June 21, 2007

Page

2

of

5

Examiner:

Group Art Unit:

OTHER (Including in this order Author, Title, Periodical Name, Pertinent Pages, Date, etc.)

| | |
|----|--|
| AR | Andersen, O., Vilsgaard Ravn, K., Sorensen, I.J., Jonson, G., Holm Nielsen, E. and Svehag, S. -E., Serum amyloid P component binds to influenza A virus haemagglutinin and inhibits the virus infection <i>in vitro</i> . <i>Scand. J. Immunol.</i> , 46:331-337, 1997. |
| BR | Ashton <i>et al.</i> , "Pentameric and decameric structures in solution of serum amyloid P component by X-ray and neutron scattering and molecular modelling analyses", <i>J. Mol. Biol.</i> 272 , 408-422, 1997. |
| CR | Askarov <i>et al.</i> , "Treatment of osteoarthritis deformans with heparin," <i>Database biosis 'Online! Biosciences information service</i> 67(3) , 190-192, 1986. |
| DR | Athanasou <i>et al.</i> , "Localized deposition of amyloid in articular cartilage," <i>Histopathology (Oxford)</i> 20(1) 41-46, 1992. |
| ER | Baltz <i>et al.</i> , "Calcium-dependent aggregation of human serum amyloid P component", <i>Biochim. Biophys. Acta</i> 701 , 229-236, 1982. |
| FR | Baltz, M.L., de Beer, F.C., Feinstein, A., Munn, E.A., Milstein, C.P., Fletcher, T.C., March, J.F., Taylor, J., Bruton, C., Clamp, J.R., Davies, A.J.S. and Pepys, M.B. Phylogenetic aspects of C-reactive protein and related proteins. <i>Ann. N. Y. Acad. Sci.</i> , 389:49-75, 1982. |
| GR | Baltz, M.L., Dyck, R.F. and Pepys, M.B., Studies of the <i>in vivo</i> synthesis and catabolism of serum amyloid P component (SAP) in the mouse. <i>Clin. Exp. Immunol.</i> , 59:235-242, 1985. |
| HR | Bartley, C.J., Orford, C.R. and Gardner, D.L., Amyloid in ageing articular cartilage. <i>J. Pathol.</i> , 145:107A, 1985. |
| IR | Booth <i>et al.</i> , "Analysis of autoaggregation and ligand binding sites of serum amyloid P component by <i>in vitro</i> mutagenesis", <i>Amyloid and Amyloidosis 1998</i> , Ed. By R.A. Kyle and M.A. Gertz, Parthenon Publishing Group, NY, 1998, pages 23-25. |
| JR | Booth <i>et al.</i> , "Instability, unfolding and aggregation of human lysozyme variants underlying amyloid fibrillogenesis", <i>Nature</i> 385 , 787-793, 1997. |
| KR | Borman, "Chemistry Highlights 2002, Medicinal and Combinatorial Chemistry," <i>Chem. Eng. News</i> 80 , 37-38, 2002. |
| LR | Breathnach, S.M., Kofler, H., Sepp, N., Ashworth, J., Woodrow, D., Pepys, M.B. and Hintner, H., Serum amyloid P component binds to cell nuclei <i>in vitro</i> and to <i>in vivo</i> deposits of extracellular chromatin in systemic lupus erythematosus. <i>J. Exp. Med.</i> , 170:1433-1438, 1989. |
| MR | Brion, P.H. and Kalunian, K.C. (2003) <i>Osteoarthritis</i> , Oxford Textbook of Medicine, 4th Ed., Vol.3 (Warrell, D.A., Cox, T.M., Firth, J.D. and Benz, E.J., Jr., eds.), Oxford University Press, Oxford, pp.62-68. |
| NR | Butler, P.J.G., Tennent, G.A. and Pepys, M.B., Pentraxin-chromatin interactions: serum amyloid P component specifically displaces H1-type histones and solubilizes native long chromatin. <i>J. Exp. Med.</i> , 172:13-18, 1990. |
| OR | Butler, P.J.G., The folding of chromatin. <i>CRC Crit. Rev. Biochem.</i> , 15:57-91, 1983. |
| PR | Cary, N.R.B., "Clinicopathological importance of deposits of amyloid in the femoral head" <i>J. Clin. Pathol.</i> , 38:868-872, 1985. |
| QR | Caspi, D., Zalzman, S., Baratz, M., Teitelbaum, Z., Yaron, M., Pras, M., Baltz, M.L. and Pepys, M.B. "Imaging of experimental amyloidosis with ¹³¹ I-serum amyloid P component." <i>Arthritis Rheum.</i> , 30:1303-1306, 1987. |
| RR | Cleaveland <i>et al.</i> , "Site of action of two novel pyrimidine biosynthesis inhibitors accurately predicted by the compare program", <i>Biochem. Pharmacol.</i> 49(7) , 947-954, 1995. |

ALL REFERENCES CONSIDERED EXCEPT WHERE LINED THROUGH. /R.L./

| | |
|-----|---|
| SR | de Beer, F.C., Baltz, M., Holford, S., Feinstein, A. and Pepys, M.B. Fibronectin and C4-binding protein are selectively bound by aggregated amyloid P component. <i>J. Exp. Med.</i> , 154:1134-1149, 1981. |
| TR | de Haas, C.J.C., van Leeuwen, E.M.M., van Bommel, T., Verhoef, J., van Kessel, K.P.M. and van Strijp, J.A.G., Serum amyloid P component bound to gram-negative bacteria prevents lipopolysaccharide-mediated classical pathway complement activation. <i>Infect. Immun.</i> , 68:1753-1759, 2000. |
| UR | Egan, M.S., Goldenberg, D.L., Cohen, A.S. and Segal, D., The association of amyloid deposits and osteoarthritis. <i>Arthritis Rheum.</i> , 25: 204-208, 1982. |
| VR | Emsley <i>et al.</i> , "Structure of pentameric human serum amyloid P component", <i>Nature</i> 367 , 338-345, 1994. |
| WR | Goffin, Y.A., Thoua, Y. and Potvliege, P.R., Microdeposition of amyloid in the joints. <i>Ann. Rheum. Dis.</i> , 40:27-33, 1981. |
| XR | Hamazaki <i>et al.</i> , "Calcium-dependent polymerization of human serum amyloid P component is inhibited by Heparin and dextran sulfate, <i>Biochimica Et Biophysica Acta</i> 998(3) , 231-235, 1989. |
| YR | Hamazaki, H., Ca ²⁺ -mediated association of human serum amyloid P component with heparan sulfate and dermatan sulfate. <i>J. Biol. Chem.</i> , 262:1456-1460, 1987. |
| ZR | Hawkins <i>et al.</i> , "Metabolic studies of radioiodinated serum amyloid P component in normal subjects and patients with systemic amyloidosis", <i>J. Clin. Invest.</i> 86 , 1862-1869, 1990. |
| AAR | Hawkins, P.N., Lavender, J.P. and Pepys, M.B., Evaluation of systemic amyloidosis by scintigraphy with ¹²³ I-labeled serum amyloid P component. <i>N. Engl. J. Med.</i> , 323:508-513, 1990. |
| BBR | Hawkins, P.N., Myers, M.J., Epenetos, A.A., Caspi, D. and Pepys, M.B., Specific localization and imaging of amyloid deposits <i>in vivo</i> using ¹²³ I-labeled serum amyloid P component. <i>J. Exp. Med.</i> , 167: 903-913, 1988. |
| CCR | Hawkins, P.N., Tennent, G.A., Woo, P. and Pepys, M.B., Studies <i>in vivo</i> and <i>in vitro</i> of serum amyloid P component in normals and in a patient with AA amyloidosis. <i>Clin. Exp. Immunol.</i> , 84:308-316, 1991. |
| DDR | Hind <i>et al.</i> , "Binding specificity of serum amyloid P component for the pyruvate acetal of galactose", <i>J. Exp. Med.</i> 159 , 1058-1069, 1984. |
| EER | Hind <i>et al.</i> , "Specific chemical dissociation of fibrillar and non-fibrillar components of amyloid deposits", <i>Lancet</i> , pages 376-378, August 18, 1984. |
| FFR | Hind, C.R.K., Collins, P.M., Baltz, M.L. and Pepys, M.B. (1985) Human serum amyloid P component, a circulating lectin with specificity for the cyclic 4,6-pyruvate acetal of galactose. Interactions with various bacteria. <i>Biochem. J.</i> 225 : 107-111. |
| GGR | Hintner, H., Booker, J., Ashworth, J., Aubock, J., Pepys, M.B. and Breathnach, S.M., Amyloid P component binds to keratin bodies in human skin and to isolated keratin filament aggregates <i>in vitro</i> . <i>J. Invest. Dermatol.</i> , 91:22-28, 1988. |
| HHR | Hohenester <i>et al.</i> , "Crystal structure of a decameric complex of human serum amyloid P component with bound dAMP", <i>J. Mol. Biol.</i> 269 , 570-578, 1997. |
| IIR | Holmgren <i>et al.</i> , "Biochemical effect of liver transplantation in two Swedish patients with familial amyloidotic polyneuropathy (FAP-met ³⁰), <i>Clin. Genet.</i> 40 , 242-246, 1991. |
| JJR | Holmgren <i>et al.</i> , "Clinical improvement and amyloid regression after liver transplantation in hereditary transthyretin amyloidosis", <i>Lancet</i> , pages 1113-1116, May 1, 1993. |
| KKR | Hutchinson <i>et al.</i> , "Human serum amyloid P component is a single uncomplexed pentamer in whole serum", <i>Mol. Med.</i> 6(6) , 482-493, 2000. |
| LLR | Hutchinson <i>et al.</i> , "The petraxins, C-reactive protein and serum amyloid P component, are cleared and catabolized by hepatocytes <i>in vivo</i> ", <i>J. Clin. Invest.</i> 94 , 1390-1396, 1994. |
| MMR | International Search Report for PCT/GB 02/03504 |
| NNR | International Search Report for PCT/GB2004/002445 |
| OOR | Iversen, "Amyloid diseases: Small drugs lead the attack," <i>Nature</i> , 414 , 231-233, 2002. |
| PPR | Kelly, "The role of glucosamine sulfate and chondroitin sulfates in the treatment of degenerative joint disease," <i>Altern. Med. Rev.</i> 3 , 27-39, 1998. |
| QQR | Klabunde <i>et al.</i> , "Rational design of potent human transthyretin amyloid disease inhibitors", <i>Nature Struct. Biol.</i> 7(4) , 312-321, 2000. |
| RRR | Ladefoged, C., Amyloid deposits in human hip joints. A macroscopic, light and polarization microscopic and electron microscopic study of congophilic substance with green dichroism in hip joints. <i>Acta Path. Microbiol. Immunol. Scand. Sect.</i> , 90:5-10, 1982. |

ALL REFERENCES CONSIDERED EXCEPT WHERE LINED THROUGH. /R.L./

| | |
|------|--|
| SSR | Ladefoged, C., Amyloid in osteoarthritic hip joints: deposits in relation to chondromatosis, pyrophosphate, and inflammatory cell infiltrate in the synovial membrane and fibrous capsule. <i>Ann. Rheum. Dis.</i> , 42: 659-664, 1983. |
| TTR | Ladefoged, C., Christensen, H.E. and Sorensen, K.H., Amyloid in osteoarthritic hip joints. Deposition in cartilage and capsule. Semiquantitative aspects. <i>Acta Orthop. Scand.</i> , 53:587-590, 1982. |
| UUR | Ladefoged, C., Amyloid deposits in the knee joint at autopsy. <i>Ann. Rheum. Dis.</i> , 45:668-672, 1986. |
| VVR | Lakhanpal, S., Li, C.Y., Gertz, M.A., Kyle, R.A. and Hunder, G.G., Synovial fluid analysis for diagnosis of amyloid arthropathy. <i>Arthritis Rheum.</i> , 30:419-423, 1987. |
| WWR | Lindorfer <i>et al.</i> , "A bispecific dsDNA monoclonal antibody construct for clearance of anti-dsDNA IGG in systemic lupus erythematosus", <i>J. Immunol. Methods</i> 248 , 125-138, 2001. |
| XXR | Liu <i>et al.</i> , "Protein heterodimerization through ligand-bridged multivalent pre-organization: enhancing ligand binding toward both protein targets," <i>J. Am. Chem. Soc.</i> 127 , 2044-2045, 2005 (published on the web Jan. 27, 2005) |
| YYR | Mitrovic, D.R., Stankovic, A., Quintero, M. and Ryckewaert, A. (1985) "Amyloid deposits in human knee and hip joints." <i>Rheumatol. Int.</i> , 5:83-89. |
| ZZR | Natchus <i>et al.</i> , "Development of new hydroxamate matrix metalloproteinase inhibitors derived from functionalized 4-aminoprolines," <i>J. Med. Chem.</i> 43 , 4948-4963, 2000. |
| AAAR | Nelson <i>et al.</i> , "Serum amyloid P component in chronic renal failure and dialysis", <i>Clinica Chimica Acta</i> 200 , 191-200, 1991. |
| BBBR | Nelson, S.R., Hawkins, P.N., Richardson, S., Lavender, J.P., Sethi, D., Gower, P.E., Pugh, C.W., Winearls, C.G., Oliver, D.O. and Pepys, M.B., Imaging of haemodialysis-associated amyloidosis with ¹²⁵ I-serum amyloid P component. <i>Lancet</i> , 338: 335-339, 1991. |
| CCCR | Nelson, S.R., Lyon, M., Gallagher, J.T., Johnson, E.A. and Pepys, M.B., Isolation and characterization of the integral glycosaminoglycan constituents of human amyloid A and monoclonal light-chain amyloid fibrils. <i>Biochem. J.</i> , 275:67-73, 1991. |
| DDDR | Noursadeghi, M., Bickerstaff, M.C.M., Gallimore, J.R., Herbert, J., Cohen, J. and Pepys, M.B. Role of serum amyloid P component in bacterial infection: protection of the host or protection of the pathogen. <i>Proc. Natl. Acad. Sci. USA</i> , 97: 14584-14589, 2000. |
| EEER | Osmand, A.P., Friedenson, B., Gewurz, H., Painter, R.H., Hofmann, T. and Shelton, E. Characterisation of C-reactive protein and the complement subcomponent C1t as homologous proteins displaying cyclic pentameric symmetry (pentaxins). <i>Proc. Natl. Acad. Sci. USA</i> , 74: 739-743, 1977. |
| FFFR | Paul <i>et al.</i> , "Identification of optimal anion spacing for anti-HIV activity in a series of cosalane tetracarboxylates", <i>Bioorg. Med. Chem. Lett.</i> 10(18) , 2149-2152, 2000. |
| GGGR | Pepys <i>et al.</i> , "Amyloid P component. A critical review.", <i>Int. J. Exp. Clin. Invest.</i> 4 , 274-295, 1997. |
| HHHR | Pepys <i>et al.</i> , "Human lysozyme gene mutations cause hereditary systemic amyloidosis", <i>Nature</i> 362 , 553-557, 1993. |
| IIIR | Pepys <i>et al.</i> , "Human serum amyloid P component is an invariant constituent of amyloid deposits and has a uniquely homogeneous glycostructure", <i>Proc. Natl. Acad. Sci. USA</i> 91 , 5602-5606, 1994. |
| JJJR | Pepys <i>et al.</i> , "Isolation of amyloid P component (Protein AP) from normal serum as a calcium-dependent binding protein", <i>Lancet</i> , pages 1029-1031, May 14, 1977. |
| KKKR | Pepys <i>et al.</i> , "Molecular mechanisms of fibrillogenesis and the protective role of amyloid P component: two possible avenues for therapy", <i>The Nature and Origin of Amyloid Fibrils (Ciba Foundation Symposium 199)</i> , John Wiley and Sons, Chichester, 1996, pages 73-89. |
| LLLR | Pepys <i>et al.</i> , "Targeted pharmacological depletion of serum amyloid P component for treatment of human amyloidosis", <i>Nature</i> 417 , 254-259, 2002. |
| MMMR | Pepys, "C-reactive protein and amyloidosis: from protein to drugs?", <i>The Lumleian Lecture, In Horizons in Medicine, Vol. 10</i> , G. Williams, editor, pp. 397-414, 1999. |
| NNNR | Pepys, M.B., Dyck, R.F., de Beer, F.C., Skinner, M. and Cohen, A.S., Binding of serum amyloid P component (SAP) by amyloid fibrils. <i>Clin. Exp. Immunol.</i> , 38:284-293, 1979. |
| OOOR | Pepys, M.B. and Baltz, M.L. Acute phase proteins with special reference to C-reactive protein and related proteins (pentaxins) and serum amyloid A protein. <i>Adv. Immunol.</i> , 34: 141-212, 1983. |
| PPPR | Pepys, M.B. and Butler, P.J.G., Serum amyloid P component is the major calcium-dependent specific DNA binding protein of the serum. <i>Biochem. Biophys. Res. Commun.</i> , 148:308-313, 1987. |
| QQQR | Pontet <i>et al.</i> , "One step preparation of both human C-reactive protein and Cit", <i>FEBS Letters</i> 88(2) , 172-175, 1978. |

ALL REFERENCES CONSIDERED EXCEPT WHERE LINED THROUGH. /R.L./

| | | |
|--|--------|---|
| | RRRR | Purkey <i>et al.</i> , "Evaluating the binding selectivity of transthyretin amyloid fibril inhibitors in blood plasma", <i>Proc. Natl. Acad. Sci. USA</i> 98(10) , 5566-5571, 2001. |
| | SSSR | Riedstra <i>et al.</i> , "Study of an anti-human transthyretin immunoadsorbent - influence of coupling chemistry on binding capacity and ligand leakage", <i>J. Chromatogr. B: Biomedical Sciences & Applications</i> . 705(2) , 213-222, 1998. |
| | TTTR | Shaker <i>et al.</i> , "Effect of low molecular weight heparin and different heparin molecular weight fractions on aggrecanase activity: structure-function relationship," <i>Blood</i> 100(11) , Abstract No. 3849, 2002. |
| | UUUR | Solomon <i>et al.</i> , Heterobifunctional Multivalent Inhibitor-Adaptor Mediates Specific Aggregates between Shiga Toxin and a Pentraxin, <i>Organic Letters</i> , American Chemical Society, 7(20) , pp. 4369-4372, 2005. |
| | VVVR | Sorensen, K.H., Stubbe Teglbjaerg, P., Ladefoged, C. and Christensen, H.E., Pyrophosphate arthritis with local amyloid deposition. <i>Acta Orthop. Scand.</i> , 52:129-133, 1981. |
| | WWWR | Srinivasan, N., Rufino, S.D., Pepys, M.B., Wood, S.P. and Blundell, T.L., A superfamily of proteins with the lectin fold. <i>Chemtracts-Biochem. Mol. Biol.</i> , 6:149-164, 1996. |
| | XXXR | Srinivasan, N., White, H.E., Emsley, J., Wood, S.P., Pepys, M.B. and Blundell, T.L. Comparative analyses of pentraxins: implications for protomer assembly and ligand binding. <i>Structure</i> , 2:1017-1027, 1994. |
| | YYR | Sukenik <i>et al.</i> , "Serum and synovial fluid levels of serum amyloid a protein and c-reactive protein in inflammatory and noninflammatory arthritis" <i>J. of Rheumatology</i> 15(6), 942-945, 1988. |
| | ZZR | Sunde, M., Serpell, L.C., Bartlam, M., Fraser, P.E., Pepys, M.B. and Blake, C.C.F., Common core structure of amyloid fibrils by synchrotron X-ray diffraction. <i>J. Mol. Biol.</i> , 273: 729-739, 1997. |
| | AAAAR | Takeda, T., Sanada, H., Ishii, M., Matsushita, M., Yamamuro, T., Shimizu, K. and Hosokawa, M., Age-associated amyloid deposition in surgically-removed herniated intervertebral discs. <i>Arthritis Rheum.</i> , 27:1063-1065, 1984.. |
| | BBBBR | Tan, S.Y., Baillod, R., Brown, E., Farrington, K., Soper, C., Percy, M., Clutterbuck, E., Madhoo, S., Pepys, M.B. and Hawkins, P.N., Clinical, radiological and serum amyloid P component scintigraphic features of β -microglobulin amyloidosis associated with continuous ambulatory peritoneal dialysis. <i>Nephrol. Dial. Transplant.</i> , 14:1467-1471, 1999. |
| | CCCCR | Tennent <i>et al.</i> , "Serum amyloid P component prevents proteolysis of the amyloid fibrils of Alzheimer disease and systemic amyloidosis", <i>Proc. Natl. Acad. Sci. USA</i> 92 , 299-4303, 1995. |
| | DDDDR | Tennent, G.A., Butler, P.J.G., Hutton, T., Woolfit, A.R., Harvey, D.J., Rademacher, T.W. and Pepys, M.B., Molecular characterization of <i>Limulus</i> polyphemus C-reactive protein. I. Subunit composition. <i>Eur. J. Biochem.</i> , 214:91-97, 1993. |
| | EEEEER | Ying, S.-C., Gewurz, A.T., Jiang, H. and Gewurz, H., Human serum amyloid P component oligomers bind and activate the classical complement pathway via residues 14-26 and 76-92 of the A chain collagen-like region of Clq. <i>J. Immunol.</i> , 150:169-176 1993. |
| | FFFFR | |
| | GGGGR | |
| | HHHHR | |
| | IIIR | |

Examiner /Ruixiang Li/

Date Considered: 02/27/2009

*EXAMINER: Initial if citation considered, whether or not citation is in conformance with MPEP § 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

ALL REFERENCES CONSIDERED EXCEPT WHERE LINED THROUGH. /R.L./